

NORTH METRO DERMATOLOGY

SKINVIVE™ CONSENT AND INSTRUCTIONS

Proposed Treatment

Injecting micro-droplets of hyaluronic acid to temporarily hydrate and smooth skin surface.

Anticipated Benefits and Limitations

Results typically last about 6 months. Results may be apparent within a few days, but full results may not be evident for 30 days. For further improvement, additional SkinVive™ can be injected 2-4 weeks following treatment. Separate charges apply for all subsequent treatments.

Possible Contraindications

- Pregnancy and nursing
- Dental work within 4 weeks of treatment
- Lidocaine allergy
- Recent injury or infection near desired treatment area

Risks

Possible side effects include: bruising, swelling, redness, pain during injection, asymmetry (unevenness), lumpiness, cold sores, numbness, granuloma or scar tissue formation, vascular occlusion, tissue necrosis, infection/abscess, and tenderness or pain after injection. The theoretical risk of unknown complications also exists, as well as lack of achieving anticipated benefits. It is important to tell your provider about all past and present medical conditions as certain conditions or medications may contraindicate treatment or increase treatment risks.

Pre-Treatment Instructions

- 10 days before treatment, avoid the following medications and supplements that may increase the risk of bruising:
Aspirin, Advil, Motrin, Ibuprofen, Aleve, Naproxen, Excedrin (all OTC pain pills except Tylenol), Vitamin E, Multi-Vitamin, Gingko Biloba, Omega-3 Fatty Acids, Cod Liver Oil, Fish Oil, Flax Oil, CoQ10, Garlic and Ginger
- If you are taking any blood-thinning medications as per doctor's orders (such as Coumadin, Warfarin, Plavix, Lovenox, Aspirin), do not discontinue without first consulting with the prescribing physician. The chance of developing an injection-related hematoma (blood clot) is higher when taking these medications.
- Do not drink alcohol within 48 hours before treatment as this may also increase the risk of bruising.
- Silagen Arnica-Bromelain Complex which is an oral supplement that can be started two days before your appointment to reduce bruising and swelling. It is also fine to start the day of the procedure, but it can be more beneficial to start a couple days in advance. This can be purchased on our website or at our clinic or by phone.
- If instructed, take prescribed medication(s) to help prevent cold sores and/or infection.

Post-Treatment Instructions

- Avoid pressure and manipulation to injected areas for 48 hours unless instructed otherwise.
- Avoid strenuous activity and exercise for 48 hours following treatment.
- Avoid exposure of the treated area to intense cold or heat for one week.
- Make-up can be worn immediately following treatment.
- It is important to contact our office immediately if pain is moderate in severity and/or if mild pain persists. You should also contact our office if observing a change in skin color or other complications. If seeking medical attention elsewhere, be sure to specify injection sites and product (this is important even if problems seem unrelated and occur several days following injections).

By signing below, I acknowledge that I have read this form and agree to the treatment with its associated risks. I hereby give consent to perform this and all subsequent Skinivive™ treatments. I hereby release North Metro Dermatology from all liabilities associated with this procedure.

Patient Name (print): _____ **Date:** _____

Patient Signature: _____

400 Village Center Drive, Suite 200 ~ North Oaks, MN 55127 ~ 651-789-9800

Rev. 11/23