NORTH METRO DERMATOLOGY Mohs Surgery/Surgical Excision Instructions

CI	П	ГΙ	ID		\Box	SI	Т	F	C	
J	<i>」</i>	ı	/ I \	_	レ	J		_	J	٠

1.	After hours, remove the outer pressure dressing. Prior to
	hours, you may replace the bandage if it becomes wet and will not
	stay adhered. Gently clean the wound once daily using a gentle soap
	and reapply antibiotic ointment (use a petrolatum-based ointment if
	allergic to antibiotic ointment). North Metro Dermatology offers an
	easy-to-use EltaMD® Dermal Wound Cleanser*.

2. Follow up for suture removal in ____ days.

NON-SUTURED OR GRANULATING WOUNDS:

1.	Keep the	dressing clear	n and dry for	hours.
----	----------	----------------	---------------	--------

- 2. Remove the dressing after _____ hours. Gently clean the wound site one or two times daily using a gentle soap (or the EltaMD® Dermal Wound Cleanser*). Apply antibiotic ointment (use a petrolatum-based ointment if allergic to antibiotic ointment) and a bandage.
- 3. Repeat wound care instructions for a minimum of _____ days or unless directed otherwise.

BLEEDING:

If bleeding occurs, apply firm pressure with a clean dressing for at least 15 minutes. Do not lift the dressing during this time to check for bleeding.

INFECTION:

Infection is uncommon after skin surgery. Some soreness, redness, and swelling are expected after surgery. Possible signs of infection include wound site becoming excessively sore, increased redness and/or wound feeling hot to the touch, excess discharge, and fever. Please contact our office if you develop any signs of infection.

BRUISING AND SWELLING:

Bruising and swelling may occur at or near the surgical site. Bruising and swelling can be especially severe around the eye resulting in a black eye and/or the eye swelling shut. Ice packs can be applied after surgery to reduce swelling.

PAIN MANAGEMENT:

ADDITIONAL INFORMATION:

Acetaminophen (Tylenol) can be taken to manage pain. Take two 325mg tablets or one or two 500mg tablet of Acetaminophen every 4-6 hours as needed. Do not take Aspirin, Ibuprofen (Advil, Motrin), Naproxen (Aleve), or other arthritis drugs as they may promote bleeding.

*This "first aid care" product is classified as an eligible expense by most HSA, HRA, and FSA plans. Check with your plan's administrator to confirm qualifying expenses, as well as any supporting documents that may be needed to substantiate your claim.

Please contact our office if you have any questions or concerns. In the event of an emergency taking place after hours, please contact Dr. Bart Endrizzi at 651-789-9819.



400 Village Center Drive, Suite 200 • North Oaks, MN 55127 • 651-789-9800 northmetroderm.com